

**10TH RESIDENTIAL COURSE ON CLINICAL
PHARMACOLOGY OF ANTIRETROVIRALS**



2005

2006

2007

2009

2010

2011

2012

2013

2014

21-22-23 January 2015

Advances in treatment of HBV

Alfredo Marzano

San Giovanni Battista Hospital

University of Turin

Italy



21-22-23 January 2015

2005
2006
2007
2009
2010
2011
2012
2013
2014

Outline

- **General background**
- **Therapy**
 - **Aims**
 - **Strategies**
 - **Efficacy**
 - **Drugs and virus**
 - **Clinical results**
 - **Summary**

10TH RESIDENTIAL COURSE ON CLINICAL
PHARMACOLOGY OF ANTIRETROVIRALS



21-22-23 January 2015

2005

2006

2007

2009

2010

2011


2012

2013

2014

Background

HBV



HBsAg+ > 400 millions
antiHBc+ > 2 billions
8 different genotypes.


High replicative activity : 10^{11-12} virions/daily
High Mutations rate: 1 every 100,000



Fig. 3. World map showing distribution of HBV genotypes. The predominant genotypes of regions of the world are shown in larger font sizes.

Natural History of CHB


Disease



1. Torresi, J, Locarnini, S. Gastroenterology. 2000.
2. Fattovich, G, Giustina, G, Schalm, SW, et al. Hepatology. 1995.
3. Moyer, LA, Mast, EE. Am J Prev Med. 1994.
4. Perrillo, R, et al. Hepatology. 2001.

Natural history of CHB

Virological aspects (infection)



Adapted by Kim HJ & Lok ASF. Hepatology 2006

HBV DNA

The Summit on
HBV Resistance

Available HBV DNA Assays

Digene Corp.


HBV Digene Hybrid-Capture I
HBV Digene Hybrid-Capture II
Ultra-Sensitive Digene Hybrid-Capture II

Roche
Molecular
Systems

Amplicor HBV Monitor
Cobas Amplicor HBV Monitor


Bayer Corp.

Versant HBV DNA 1.0
Versant HBV DNA 3.0




qHBsAg

117 chronic hepatitis B patients
followed for 99 ± 16 months; number of visits 17 (8–49)



Staging

Liver biopsy and Fibroscan



Gaia, J Hepatol 2011

* Oliveri, WJG 2008

10TH RESIDENTIAL COURSE ON CLINICAL
PHARMACOLOGY OF ANTIRETROVIRALS

2005

2006

2007

2009

2010

2011

2012

2013

2014



21-22-23 January 2015

Therapy aims

Antiviral Therapy in HBV Goals



		HBV DNA IU	qHBsAg IU	HBeAg
AC		> 2000	> 1000	Pos/neg
IC		< 2000	<1000	neg
		neg	<100	neg
OBI		neg	neg	neg

STOP

Evolution

Cirrhosis (regression)

Portal hypertension

Liver transplantation

HCC

Mortality

10TH RESIDENTIAL COURSE ON CLINICAL
PHARMACOLOGY OF ANTIRETROVIRALS

2005

2006

2007

2009

2010

2011

2012

2013

2014



21-22-23 January 2015

Therapy Strategies

Therapeutic strategies

PEG
NUC (e+)

Curative (defined time)

Suppressive (NUC) (undefined time)

This topic falls outside my time

10TH RESIDENTIAL COURSE ON CLINICAL
PHARMACOLOGY OF ANTIRETROVIRALS

2005

2006

2007

2009

2010

2011

2012

2013

2014



21-22-23 January 2015


Therapy

Efficacy




PEG-IFN

Peg-IFN in HBeAg-positive




J Hepatology 2012

50% HBsAg clearance



Peg-IFN alpha-2a in HBeAg-negative CHB. Sustained response (5 yr)





Marcellin et al, APAASL 2009

Better response in genotype non-D/E

Marcellin, Gastroenterology 2009


ETV

TDF



Marengo & Marzano
Antivir Ther 2013

Marengo & Marzano
APT 2014



Pangenotypic



21-22-23 January 2015

Therapy

Drugs and Virus


Why the mono-therapy is effective?

Virus and Drugs characteristics

Suppression vs. cure: viral biology is the basis

HBV

10 genotypes
(Latent Reservoir)




Long-time cccDNA
“Low” replicative space

Suppression

HCV

6 genotypes
(No Latent Reservoir)



**Definitive Viral
Clearance**

Cure


Drugs

Pharmacologic barrier

Intracellular drugs levels and resistance

I generation (high drug levels)

Low toxicity, medium potency




Drugs

Pharmacologic barrier

Intracellular drugs levels and resistance

II generation (low drug levels)

High toxicity, medium potency




Drugs

Pharmacologic barrier


Intracellular drugs levels and resistance

III generation (high drug levels)

Low toxicity, high potency



Drug levels, Genetic barrier and the virus




HBV

HCV

Virus	HBV	HCV
daily production of virions per day	$10^{12} - 10^{13}$	10^{12}
half-life of free virions (h)	3-24	2-3
half-life of intracellular virions	months (dependent on infected cells $t_{1/2}$)	hours (not dependent on infected cells $t_{1/2}$)
mutation rate	high	very high
constraints due to ORFs in targeted viral enzymes	high	none
immune-mediated escape mutants	infrequent	frequent
Target cells		
half-life of infected cells	months	weeks
size of susceptible cells compartment	small	probably large
intracellular viral reservoir	yes (integrated cDNA)	no
	yes (cccDNA)	

Genetic Barrier

Incidence of Resistance in NUC-naïve Patients





21-22-23 January 2015

2005
2006
2007
2009
2010
2011
2012
2013
2014



Therapy

Clinical Results


**Can we reach clinical results
with the NUCs?**

Regression of histological cirrhosis with TDF

Marcellin P; The Lancet 2013




Reversal of fibrosis/cirrhosis with ETV



* Median time of long-term biopsy: 6 years (range: 3–7 years)

Clinical experience with III gen NUCs:



Entecavir 100 pts (55 compensated cirrhosis)



— Cirrhosis
 — CH

Changes of esophageal varices (EV) in compensated cirrhotics treated with LAM±TDF for 10 years

Overall, EV worsening rate per year: 0.9%*



* 6 of 7 progressors (86%) had either LMV-R and/or HCC

Decompensated cirrhosis

Table 1 Studies of nucleos/tide analogues in patients with HBV-induced decompensated cirrhosis

Study(reference)	Fontana ²⁶	Schiff ²⁷	Liaw* ³⁰		Liaw ²⁸		Shim ²⁹	
No. of patients treated	154	226	45	45	22	100	91	70
Drug(s) used	LAM	ADV+LAM	TDF	TDF+FTC	ETV	ETV	ADV	ETV
Baseline characteristics								
Lamivudine experienced	0%	100%	42%	38%	36%	36%	33%	0%
HBV DNA (log ₁₀ c/ml)	~7	7	5.7	6.3	5.9	7.5	8.2	7.2
CP score	9	60% B/C	7	7	7	8.8	8.4	8.4
MELD score	NR	NR	11.0	13.0	10.5	17.1	15.3	11.5
1-year efficacy and safety data								
Undetectable HBV DNA	NR	59%	71%	88%	73%	57%	20%	89%
CP \geq 2 point decrease	NR	~50%	26%	48%	42%	35%	27%	49%
Decrease in MELD score	NR	2	2	2	2	2.6	1.7	2.3
Mortality	16%	14%	4%	4%	9%	12%†	12%†	12.9%§
HCC	NR	NR	NR	NR	NR	12%‡	20%‡	6.9%
Cr \geq 0.5 mg/dl increase	NR	6%	9%	2%	5%	17%‡	24%‡	NR
Phosphorus < 2.0 mg/dl	NR	NR	2%	4%	3%	NR	NR	NR
Drug resistance	27%	0	0%	0%	0%	0%	0%	NR

*Patients with 2 log decrease in HBV DNA at week 8 can switch to open label TDF + FTC.

†Mortality at 24 weeks, cumulative rates of mortality were 23% and 33% and of LT 11% and 3%, respectively, for ETV and ADV.

‡Cumulative rates.

§Mortality or liver transplant.

ADV, adefovir dipivoxil; CP, Child–Pugh score; ETV, entecavir; HBV, hepatitis B virus; HCC, hepatocellular carcinoma; LAM, lamivudine; TDF, tenofovir disoproxil fumarate. FTC, emtricitabine; MELD, model for end-stage liver disease; NR, not recorded; MELD; model for end-stage liver disease.

Prophylaxis of hepatitis B recurrence after LT

2429 pts




Figure 1: Subgroups of patients under antiviral prophylaxis against HBV recurrence after liver transplantation: 1910 (79%) patients received HBIG and LAM, 304 (12.5%) HBIG and ETV/TDF, 103 (4%) ETV/TDF after HBIG discontinuation and 112 (4.5%) ETV/TDF without any HBIG. HBIG= hepatitis B immunoglobulin; LAM= lamivudine; ETV= entecavir; TDF= tenofovir.






Figure 2: Risk of recurrence hepatitis B virus (HBV) infection after liver transplantation (LT) in relation to the type of posttransplant HBV prophylaxis. HBIG = hepatitis B immunoglobulin; LAM = lamivudine; ETV = entecavir; TDF = tenofovir.


Excellent results but.....

HCC remains the residual problem




Expected annual risk in Europe (untreated) 0.2% (CH) -2% (Cirrhosis)

HCC rates in NUC-naïve cirrhotic patients long-term responding to NUC (LAM-exp)




Num. of pts:	211	81	62	42
Drug:	LAM	LAM	LAM	LAM
Study:	RCT	Review	Retrospect.	Retrospect.
Follow-up:	3 yrs	2 yrs	6 yrs	5 yrs

Clinical experience with ETV: HCC in 472 (gen B-C) vs 1143 untreated patients



HCC rates in ETV or TDF treated CHB pts European multicenter study

(N=1231 patients, 54% naive, 55% TDF, 39 months follow-up)



LT for HBV-related cirrhosis in US and Europe

Van Bommel, Liver Int 2013; Burra & Marzano, J Hepatol 2013







Fig. 1. Number of patients on the waiting list for liver transplantation for hepatitis B-related indications in the United States (30).



* $p < 0.001$ vs. 1988-1995

Fig. 1. Evolution of liver transplantation for virus-related chronic liver disease in Europe according to different time periods. (A) Evolution of liver transplantation for HBV-related liver disease considering the whole cohort of study. (B) Evolution of liver transplantation for HBV-related liver disease according to the indication for liver transplantation (HBVdec and HBV/HCC).



$\dagger p = 0.87$ vs. 1988-1995;
* $p < 0.001$ vs. 1988-1995

Summary



DE BEERS
A DIAMOND IS FOREVER



HBV too

