

**10TH RESIDENTIAL COURSE ON CLINICAL
PHARMACOLOGY OF ANTIRETROVIRALS**



21-22-23 January 2015

Starhotels Majestic
corso Vittorio Emanuele II 54 - **TURIN**



2005

2006

2007

2009

2010

2011

2012

2013

2014

2015

10TH
EDITION

Drug drug interaction of new HCV drugs

Stefano Bonora

University of Torino


Disposition of “old” DAAs

Drug	Dosing regimen	CYP	P-glycoprotein	Non-CYP metabolism
Telaprevir	Q8h No significant boosting by RTV	CYP 3A4: ▪ Substrate ▪ Inhibitor	▪ Substrate ▪ Inhibitor	—
Boceprevir	tid No significant boosting by RTV	CYP 3A4/5: ▪ Substrate ▪ Inhibitor	▪ Substrate	AKR ▪ Substrate


	VICTIM of DDI	PERPETRATOR of DDI	DDI potential
Teleprevir	Substrate for CYP 3A4, PgP	Inhibits CYP 3A4, PgP, OATP1B1/2 ? Protein binding	Significant
Boceprevir	Substrate for aldoketoreductase, CYP 3A4, PgP, BCRP	Inhibits CYP 3A4, PgP, OCT 1&2	Significant
Simeprevir	Substrate for CYP 3A4, PgP	Inhibits OATP1B1, MRP2 Mild inhibitor gut CYP 3A4, PgP	Moderate
Sofosbuvir	cathepsin A, esterases, kinases PgP & BCRP substrate (parent)	Weak inhibitor of gut PgP & BCRP	Low
Ledipasvir	Primarily excreted unchanged (>98% faeces), PgP / BCRP substrate	Weak inhibitor of PgP/BCRP, ?OATP1B1/3	?
ABT450r	Substrate for CYP 3A4, PgP, OATP1B1/3	Weak inhibitor PgP/BCRP (gut), ?OATP1B1/3	Moderate to Significant (RTV)
Ombitasvir (ABT-267)	Substrate for PgP, BCRP (CYP 3A4)	Weak inhibitor of UGT1A1	
Dasabuvir (ABT-333)	Substrate of CYP 2C8 > 3A4 > 2D6, Substrate of PgP, BCRP	Weak inhibitor of UGT1A1	
Daclatasvir	Substrate for CYP 3A4, PgP	Inhibits OATP1B1/3 & PgP	Moderate
Asuneprevir	Substrate for OATP1B1/2B1 CYP 3A4	Inhibits CYP2D6 (mod) & OATP1B1/3 (weak), ?BCRP, Weak CYP3A4 inducer	?
Faldeprevir	Substrate for CYP 3A4, PgP, OATP1B1 & MRP2	Inhibitor of CYP 3A4, 2C9, UGT1A1, Probably inhibits OATP1B1/3, MRP2	Moderate
MK-5172	Substrate for CYP 3A4, PgP, ? OATP1B1	Inhibits CYP 2C8, weak inhibitor of UGT1A1, ? BCRP	Moderate
MK-8742	Substrate for CYP 3A4, PgP, ?OATP1B1	weak inhibitor of UGT1A1	Moderate

	VICTIM of DDI	PERPETRATOR of DDI	DDI potential
Teleprevir	Substrate for CYP 3A4, PgP	Inhibits CYP 3A4, PgP, OATP1B1/2 ? Protein binding	Significant
Boceprevir	Substrate for aldoketoreductase, CYP 3A4, PgP, BCRP	Inhibits CYP 3A4, PgP, OCT 1&2	Significant
Simeprevir	Substrate for CYP 3A4, PgP	Inhibits OATP1B1, MRP2 Mild inhibitor gut CYP 3A4, PgP	Moderate
Sofosbuvir	cathepsin A, esterases, kinases PgP & BCRP substrate (parent)	Weak inhibitor of gut PgP & BCRP	Low
Ledipasvir	Primarily excreted unchanged (>98% faeces), PgP / BCRP substrate	Weak inhibitor of PgP/BCRP, ?OATP1B1/3	?
ABT450r	Substrate for CYP 3A4, PgP, OATP1B1/3	Weak inhibitor PgP/BCRP (gut), ?OATP1B1/3	Moderate to Significant (RTV)
Ombitasvir (ABT-267)	Substrate for PgP, BCRP (CYP 3A4)	Weak inhibitor of UGT1A1	
Dasabuvir (ABT-333)	Substrate of CYP 2C8 > 3A4 > 2D6, Substrate of PgP, BCRP	Weak inhibitor of UGT1A1	
Daclatasvir	Substrate for CYP 3A4, PgP	Inhibits OATP1B1/3 & PgP	Moderate
Asuneprevir	Substrate for OATP1B1/2B1 CYP 3A4	Inhibits CYP2D6 (mod) & OATP1B1/3 (weak), ?BCRP, Weak CYP3A4 inducer	?
Faldeprevir	Substrate for CYP 3A4, PgP, OATP1B1 & MRP2	Inhibitor of CYP 3A4, 2C9, UGT1A1, Probably inhibits OATP1B1/3, MRP2	Moderate
MK-5172	Substrate for CYP 3A4, PgP, ? OATP1B1	Inhibits CYP 2C8, weak inhibitor of UGT1A1, ? BCRP	Moderate
MK-8742	Substrate for CYP 3A4, PgP, ?OATP1B1	weak inhibitor of UGT1A1	Moderate

Mechanisms of drug-drug interactions involving DAAs



Drug transporters




DAA	Substrate	Effect	Clinical DDI extent	Comments
Simeprevir	CYP3A4 P-gp	Inhibits OATP1B1 and multi-drug resistant protein 2 Inhibits gut CYP3A4 and P-gp	Moderate	Bile elimination
Paritaprevir/r	CYP3A4 P-gp and OATP1B1/3	Inhibits liver CYP3A4 Inhibits gut P-gp, BCRP Inhibits OATP1B1/3	Significant	Efavirenz and rilpivirine contraindicated Hyperbilirubinemia
Sofosbuvir	Cathepsin A, esterases and kinases P-gp and BCRP	Inhibits gut P-gp	<u>Low</u>	Urine elimination (80%) and bile (14%)
Dasabuvir	CYP2C8, 3A4 and 2D6 P-gp and BCRP	Inhibits UGT1A1 Inhibits OATP1B1	Significant	
Daclatasvir	CYP3A4 and P-gp	Inhibits OATP1B1/3 and P-gp	Moderate	Increases statin exposure Darunavir/r increases daclatasvir exposure
Ledipasvir	P-gp and BCRP	Inhibits gut P-gp and BCRP Inhibits OATP1B1/3	Low	Increases statin exposure
Ombitasvir	CYP3A4 P-gp and BCRP	Inhibits CYP2C8 and UGT1A1	Significant	

BCRP: Breast cancer resistance protein; DDA: Direct-acting antivirals; DDI: Drug-drug interactions; MRP: Multidrug resistance protein; OATP: Organ anion transporter protein; P-gp: P glycoprotein; UGT: Uridin-glucoronil-transferase.

Sofosbuvir Intracellular Activation

- As nucleotide analog prodrug
SOF is activated by sequential metabolic pathways including
 - Low affinity and high capacity hydrolases (CES1, CatA and HINT1)
 - Nucleotide phosphorylation (UMP-CMP kinase and NDP kinase)
- Only SOF can enter hepatocytes and be converted to active TP (GS-461203)
- SOF can also undergo extrahepatic metabolism to form GS-331007 (predominant metabolite) principally eliminated in urine



Effect of ARVs on Sofosbuvir: *Victim*

Drug	Effect on Sofosbuvir and GS-331007 AUC (exposure)	Recommendation
Darunavir/r	SOF increased 34%; GS-331007 – no effect	No dose adjustment
Rilpivirine	No effect on SOF or GS-331007	No dose adjustment
Efavirenz	No effect on SOF or GS-331007	No dose adjustment
Raltegravir	No effect on SOF or GS-331007: RAL decreased 27%	No dose adjustment
Tenofovir	No effect on SOF or GS-331007	No dose adjustment

Table 15. Potentially Significant Drug Interaction

Concomitant Drug Class: Drug Name	Clinical Comment
<u>Anti-convulsants:</u> Carbamazepine Phenytoin Phenobarbital Oxcarbazepine	Co-administration of SOF with carbamazepine, phenytoin, phenobarbital, or oxcarbazepine is expected to decrease the concentration of SOF, due to potential induction of Pgp. This may result in a reduced therapeutic effect. Such co-administration is not recommended.
<u>Anti-mycobacterials:</u> Rifampin Rifabutin Rifapentine	<p>Co-administration of SOF with rifabutin or rifapentine is expected to decrease the concentration of SOF, due to potential induction of Pgp. This may result in a reduced therapeutic effect. Such co-administration is not recommended.</p> <p>SOF should not be used with rifampin, a potent intestinal Pgp inducer.</p>
<u>Anti-retrovirals:</u> Tipranavir/ritonavir	Co-administration of SOF with ritonavir-boosted tipranavir is expected to decrease the concentration of SOF, due to potential induction of Pgp. This may result in a reduced therapeutic effect. Such co-administration is not recommended.
<u>Herbal Supplements:</u> St John's Wort	SOF should not be used with St John's Wort, a potent intestinal Pgp inducer.

↓ = decrease; Pgp = p-glycoprotein; SOF = sofosbuvir

DAAs in Development

Drug	CYP Activity	Transporters	Interaction Potential
Ledipasvir	<ul style="list-style-type: none"> ▪ Little metabolism ▪ Not Inhibitor of CYP or UGT ▪ Not Inducer of CYP or UGT 	<ul style="list-style-type: none"> ▪ P-gp substrate (likely) ▪ Inhibition of intestinal P-gp (weak) ▪ Inhibition of OATP1B1/3 (weak) 	<ul style="list-style-type: none"> ▪ Weak

Eley T et al, 2013, 8th Int Workshop on Clin Pharm of Hep Ther; Abs O-13; Eley T et al, 2011, 62nd AASLD Abs 381; Eley T et al 2012, 7th Int Workshop on Clin Pharm of Hep Ther; Abs O-4; Kirby B et al 2013, 8th Int Workshop on Clin Pharm of Hep Ther; Abs O-20; Mathias A, 14th Int Workshop on Clin Pharm of HIV Ther, Session 5

Effect of P-gp Inducers on LDV/SOF²

Object	Perpetrator	AUC	C _{max}
SOF	Rifampin	↓ 72%	↓ 77%
GS-331007		↔	↔
LDV		↓ 56%	↓ 35%

- ♦ P-gp inducers (eg, rifampin, St. John's wort) should not be used with LDV/SOF
- ♦ Use of other P-gp inducers (eg, rifabutin, rifapentine, phenytoin, phenobarbital, carbamazepine, and oxcarbazepine) with LDV/SOF is not recommended



Effect of LDV/SOF on OATP Substrates⁴

Object	Perpetrator	AUC	C _{max}
Pravastatin	LDV*	↑ 168%	↑ 166%
Rosuvastatin		↑ 699%	↑ 1670%

*LDV administered in combination with VDV and TGV.

- ♦ VDV is a potent OATP inhibitor; LDV is a weak OATP inhibitor
- ♦ SOF and GS-331007 are not OATP inhibitors
- ♦ LDV/SOF may be administered with OATP substrates
- ♦ Overall incidence of statin-related adverse events (eg, myopathy, fatigue, asthenia) in the pooled Phase 2/3 population was similar in HCV-infected patients who did and did not receive statins
- ♦ Clinically relevant interactions are not expected with LDV/SOF and most statins (eg, pravastatin); the use of rosuvastatin is not recommended

Effect of LDV/SOF on TFV (ATR or CPA)



TFV PK Parameter	Mean (%CV)		GMR% (90% CI): LDV/SOF + ARV vs ARV	
	LDV/SOF + ATR	LDV/SOF + CPA	ATR	CPA
AUC _{tau} (ng·h/mL)	4400 (27.1)	4780 (28.6)	198 (177, 223)	140 (131, 150)
C _{max} (ng/mL)	527 (29.9)	490 (24.1)	179 (156, 204)	132 (125, 139)
C _{tau} (ng/mL)	113 (33.0)	118 (26.4)	263 (237, 297)	191 (174, 210)

Data presented to 3 significant figures; LDV/SOF + ATR: n=15; ATR: n=17; LDV/SOF + CPA: n=14; CPA: n=14.


◆ LDV/SOF increased TFV exposure

- Similar **absolute** TFV exposures (within ATR or CPA) with LDV/SOF
 - TFV exposures in LDV/SOF + NNRTI-based regimens are similar to those with boosted HIV PIs
- Lack of marked changes in TFV Cl_{renal} (data on file)


DAA	Substrate	Effect	Clinical DDI extent	Comments
Simeprevir	<u>CYP3A4</u> P-gp	Inhibits <u>OATP1B1</u> and multi-drug resistant protein 2 Inhibits gut <u>CYP3A4</u> and P-gp	Moderate	Bile elimination
Paritaprevir/r	CYP3A4 P-gp and OATP1B1/3	Inhibits liver CYP3A4 Inhibits gut P-gp, BCRP Inhibits OATP1B1/3	Significant	Efavirenz and rilpivirine contraindicated Hyperbilirubinemia
Sofosbuvir	Cathepsin A, esterases and kinases P-gp and BCRP	Inhibits gut P-gp	Low	Urine elimination (80%) and bile (14%)
Dasabuvir	CYP2C8, 3A4 and 2D6 P-gp and BCRP	Inhibits UGT1A1 Inhibits OATP1B1	Significant	
Daclatasvir	CYP3A4 and P-gp	Inhibits OATP1B1/3 and P-gp	Moderate	Increases statin exposure Darunavir/r increases daclatasvir exposure
Ledipasvir	P-gp and BCRP	Inhibits gut P-gp and BCRP Inhibits OATP1B1/3	Low	Increases statin exposure
Ombitasvir	CYP3A4 P-gp and BCRP	Inhibits CYP2C8 and UGT1A1	Significant	

BCRP: Breast cancer resistance protein; DDA: Direct-acting antivirals; DDI: Drug-drug interactions; MRP: Multidrug resistance protein; OATP: Organ anion transporter protein; P-gp: P glycoprotein; UGT: Uridin-glucoronil-transferase.


Simeprevir Exhibits Non-Linear Pharmacokinetics




Trial C101 (healthy subjects)



No Correlation Between Efficacy (SVR12) and Exposures Achieved with 150 mg QD




An Increased Incidence of Rash was Associated with Higher Exposures in Phase 3



Similar relationships between exposures and:

- photosensitivity
- pruritus
- dyspnea
- increased bilirubin


Systemic Exposures were Higher in Asian Patients in Phase 3 trials




Relative to Caucasian subjects,
Asian subjects have:

smaller liver
less CYP3A
less OATP1B1

slower clearance
higher exposure



HCV-Uninfected Subjects with Moderate or Severe Hepatic Impairment had Higher Exposures Compared to Healthy Controls



- PegIFN is contraindicated in Child-Pugh class B or C
- Simeprevir PK will be evaluated in patients with moderate or severe hepatic impairment during ongoing IFN-free development

Effect of ARVs on Simeprevir: *Victim*

Drug	Effect on Simeprevir AUC (exposure)	Mechanism/ <i>Recommendation</i>
Darunavir/r	2.6-fold increase (DRV increased 18%)	RTV Inhibits CYP3A4 <i>Not recommended</i>
Rilpivirine	No effect	<i>No dose adjustment</i>
Efavirenz	70% decrease	EFV induces CYP3A4 <i>Not recommended</i>
Raltegravir	11% decrease	<i>No dose adjustment</i>
Tenofovir	14% decrease (TFV increased 18%)	Intestine or renal transport <i>No dose adjustment</i>

Figure 6: Effect of SMV Administration at 150 mg Once Daily on Exposure of Coadministered Drugs

HIV antiretrovirals

Ritonavir¹
 Darunavir¹
 Rilpivirine
 Tenofovir Disoproxil Fumarate
 Efavirenz
 Raltegravir

Immunosuppressants

Tacrolimus
 Cyclosporine

HMG-CoA reductase

Inhibitors


Atorvastatin
 Simvastatin
 Rosuvastatin

Oral contraceptives

Ethinylestradiol
 Norethindrone

Other drugs

Midazolam (oral)
 Midazolam (iv)
 S-Warfarin
 Caffeine
 Omeprazole
 Dextromethorphan
 Escitalopram
 R (-) Methadone
 S (+) Methadone
 Digoxin
 Erythromycin
 Rifampin²



Prevalence of Drug-Drug Interactions upon Addition of Simeprevir- or Sofosbuvir-Containing Treatment to Medication Profiles of Patients with HIV and Hepatitis C Coinfection.

Patel N et al. AIDS Res Human Retrovir 2014

- DDIs were present in 20% of the 335 included patients.
- After the addition of **SIM-containing therapy**, the frequency of DDIs significantly **increased to 88.4%** ($p<0.001$).
- After adding **SOF-containing therapy**, the prevalence of DDIs increased to 24.5% ($p<0.001$).
- Variables independently associated with DDIs after the addition of **SIM-containing therapy** were **NNRTI regimen** (prevalence ratio, PR: 1.62; 95% confidence interval, CI: 1.38-1.91, $p<0.001$), **PI regimen** (PR: 1.64; 95% CI: 1.40-1.93, $p<0.001$), and **≥ 7 non-HIV medications** (PR: 1.06; 95% CI: 1.00-1.14, $p=0.09$).
- *The prevalence of DDIs was significantly lower for SOF-containing HCV therapy within various types of ART regimens.*

Worthy to get PK data in real life?

- ✓ Simeprevir:
 - No DDI data with DTG, ATV/r, LPV/r, COBI, MVC
 - overdosing in cirrhotic patients and/or HIV coinfectd treated with PI/r
 - increase risk of rash/photosensitivity?
 - dose decrease in cirrhotics?

DAA	Substrate	Effect	Clinical DDI extent	Comments
Simeprevir	CYP3A4 P-gp	Inhibits OATP1B1 and multi-drug resistant protein 2 Inhibits gut CYP3A4 and P-gp	Moderate	Bile elimination
Paritaprevir/r	CYP3A4 P-gp and OATP1B1/3	Inhibits liver CYP3A4 Inhibits gut P-gp, BCRP Inhibits OATP1B1/3	Significant	Efavirenz and rilpivirine contraindicated Hyperbilirubinemia
Sofosbuvir	Cathepsin A, esterases and kinases P-gp and BCRP	Inhibits gut P-gp	Low	Urine elimination (80%) and bile (14%)
Dasabuvir	CYP2C8, 3A4 and 2D6 P-gp and BCRP	Inhibits UGT1A1 Inhibits OATP1B1	Significant	
<u>Daclatasvir</u>	<u>CYP3A4 and P-gp</u>	<u>Inhibits OATP1B1/3 and P-gp</u>	Moderate	Increases statin exposure Darunavir/r increases daclatasvir exposure
Ledipasvir	P-gp and BCRP	Inhibits gut P-gp and BCRP Inhibits OATP1B1/3	Low	Increases statin exposure
Ombitasvir	CYP3A4 P-gp and BCRP	Inhibits CYP2C8 and UGT1A1	Significant	

BCRP: Breast cancer resistance protein; DDA: Direct-acting antivirals; DDI: Drug-drug interactions; MRP: Multidrug resistance protein; OATP: Organ anion transporter protein; P-gp: P glycoprotein; UGT: Uridin-glucuronil-transferase.

Daclatasvir DDIs - *victim*

CYP3A4 and/or Pgp INDUCERS - anti-HIV	EFV	Increase to 90 mg/day
CYP3A4 and/or Pgp INDUCERS - other than EFV	<i>Etravirine</i> , Carbamazepine, oxacarbazepine, phenobarbital, dextame St John's wort	Not recommended
CYP3A4 and/or Pgp INHIBITORS Anti-HIV	ATV/RTV DRV/r and LPV/R	Decrease to 30 mg/day Standard dose*
CYP3A4 and/or Pgp INHIBITORS other than anti-HIV	clarithromycin, itraconazole, quinidine, ranolazine	Caution or decrease to 30 mg/day

* Daclatasvir AUC increase by 40% (DRV/r) and 15% (LPV/R)- HEP DART meeting Dec 2014

Daclatasvir DDIs - *perpetrator*

- ✓ No effect of gastric acid modifiers, midazolam or oral contraceptives
- ✓ Caution with rosuvastatin (increase of AUC by 58%%)

Worthy to get PK data in real life?

✓ **Simeprevir:**

- No DDI data with DTG, ATV/r, LPV/r, COBI, MVC
- overdosing in cirrhotic patients and/or HIV coinfectd treated with PI/r
 - increase risk of rash/photosensitivity?
 - dose decrease in cirrhotics?

✓ **Daclatasvir**

- No DDIs data with ETV, DTG, COBI, MVC
- HIV-coinfectd with dose reduction (ATV, COBI): underdosing?

DDIs of daclatasvir have been evaluated in healthy volunteers

PK exposure was shown to be **lower in HCV+** as compared to healthy volunteers

Pk exposure was shown to be **lower in cirrhotics patients (HCV-)** as compared to healthy volunteers (but unbound fraction equal)

+

+

+

=

- HIV/HCV cirrhotic,
- ATV/r or COBI-containing HAART
- Daclatasvir dose reduction (30 mg OD)

Magnitude of DDI and drug exposure are not easy to predict

Journal of Antimicrobial Chemotherapy Advance Access published October 29, 2013

J Antimicrob Chemother
doi:10.1093/jac/dkt423

**Journal of
Antimicrobial
Chemotherapy**

Modelling clinical data shows active tissue concentration of daclatasvir is 10-fold lower than its plasma concentration

Ruian Ke^{1*}, Claude Loverdo¹, Hangfei Qi², C. Anders Olson², Nicholas C. Wu³, Ren Sun²⁻⁴
and James O. Lloyd-Smith^{1,5}

DAAs in Development

Drug	CYP Activity	Transporters	Interaction Potential
Asunaprevir	<ul style="list-style-type: none">▪ CYP3A4 substrate▪ Inducer of CYP3A4 (weak)▪ Inhibition of CYP2D6 (weak)	<ul style="list-style-type: none">▪ P-gp, OATP1B1/3 substrate▪ Inhibition of P-gp (weak), OATP1B1/3	<ul style="list-style-type: none">▪ Moderate


Eley T et al, 2013, 8th Int Workshop on Clin Pharm of Hep Ther; Abs O-13; Eley T et al, 2011, 62nd AASLD Abs 381; Eley T et al 2012, 7th Int Workshop on Clin Pharm of Hep Ther; Abs O-4; Kirby B et al 2013, 8th Int Workshop on Clin Pharm of Hep Ther; Abs O-20; Mathias A, 14th Int Workshop on Clin Pharm of HIV Ther, Session 5

DRUG INTERACTIONS WITH ABBVIE'S 3D REGIMEN (ABT450/RITONAVIR, OMBITASVIR AND DASABUVIR)

	ABT-450/r	Ombitasvir (ABT-267)	Dasabuvir (ABT-333)
Pharmacology	NS3/4A protease inhibitor	NS5A inhibitor	NS5B non-nucleoside inhibitor
Adult Dose	<i>Investigational:</i> 150 mg once daily with ritonavir 100 mg once daily Co-formulated with ritonavir and ombitasvir as 150/100/25 mg tablet.	<i>Investigational:</i> 25 mg once daily Coformulated with ABT-450 and ritonavir (150 mg ABT450, 100 mg ritonavir, 25 mg ombitasvir fixed dose tablet).	<i>Investigational:</i> 400 mg BID
Impact of Food	In healthy subjects, ABT- 450 C _{max} , and AUC were 11 to 19% higher under non-fasting conditions compared to fasting. ABT- 450/r can be given with or without food ¹		
Kinetic Characteristics	Substrate of 3A4, P-gp, OATP1B1. Inhibits CYP2C8, UGT1A1, OATP1B1 and OATP1B3	Substrate of 3A4, P-gp. Inhibits CYP2C8, UGT1A1.	Substrate of CYP2C8>3A4, 2D6, P-gp. Inhibits UGT1A1, OATP1B1.



Why is ABT-450 Dosed with Ritonavir?



- Significant pharmacokinetic boosting allows for QD administration at lower ABT-450 doses while potentially improving the resistance profile.
- Changes in ABT-267 and ABT-333 exposures were $\leq \sim 50\%$ when dosed with ABT-450 + ritonavir.

DDI Study of 3D Regimen with	Effect of HIV PI on C_{max} and AUC of 3D Regimen	Effect of 3D Regimen on C_{max} , AUC and C_{trough} of HIV PI
ATV + RTV QD ✓	ABT-450 $\leq 94\%$ ↑ ; ombitasvir $\leq 23\%$ ↓ ; dasabuvir ↔	ATV ↔
DRV + RTV QD ✓	ABT-450 $\leq 54\%$ ↑; ombitasvir ↔ ; dasabuvir ↔	DRV C_{max} and AUC $\leq 24\%$ ↓ and C_{trough} 48% ↓
DRV + RTV BID ✓	ABT-450 $\leq 41\%$ ↓; ombitasvir $\leq 27\%$ ↓ ; dasabuvir $\leq 27\%$ ↓	DRV C_{max} and AUC ↔ and C_{trough} 43% ↓
LPV/r BID ✗	ABT-450 $\leq 117\%$ ↑ ; ombitasvir ↔ ; dasabuvir ↔	LPV ↔

DDI Study of 3D Regimen with	Effect of HIV-1 ARV Drugs on C_{max} and AUC of 3D Regimen	Effect of 3D regimen on C_{max} , AUC and C_{trough} HIV-1 ARV Drugs
FTC + TDF ✓	ABT-450 $\leq 32\%$ ↓; ombitasvir ↔ ; dasabuvir ↔	FTC ↔; TDF C_{max} and AUC ↔ , C_{trough} 24% ↑
RAL ✓	ABT-450 ↔ ; ombitasvir ↔ ; dasabuvir ↔	RAL $\leq 134\%$ ↑
RPV ✗	ABT-450 $\leq 30\%$ ↑ ; ombitasvir ↔ ; dasabuvir ↔	RPV C_{max} 155% ↑ , AUC 225% ↑ and C_{trough} 262% ↑

MK-5172 e MK-8742

DRUG INTERACTIONS WITH MK-5172 AND MK-8742

	MK-5172 (Merck)	MK-8742 (Merck)
Pharmacology	NS3/4A protease inhibitor	NS5A inhibitor
Adult Dose	<i>Investigational:</i> 100 mg once daily	<i>Investigational:</i> 50 mg once daily
Kinetic Characteristics	<div>Substrate of CYP3A4, P-gp and OATP1B1.¹</div> <div>Inhibitor of CYP2C8, 3A4 (weak), UGT1A1 (weak) and possibly BCRP.</div>	<div>Substrate of CYP3A4, P-glycoprotein (P-gp) and the organic anion-transporting</div> <div>polypeptide (OATP) in vitro. No age effect observed in young (22-45 yrs) vs elderly (65-78 yrs) males; ~33% higher AUC in elderly female vs male subjects after adjusting for body weight.²</div>

Worthy to get PK data in real life?

✓ **Simeprevir:**

- No DDI data with DTG, ATV/r, LPV/r, COBI, MVC
- overdosing in cirrhotic patients and/or HIV coinfecting treated with PI/r
 - increase risk of rash/photosensitivity?
 - dose decrease in cirrhotics?

✓ **Daclatasvir**

- No DDIs data with ETV, DTG, COBI, MVC
- HIV-coinfecting with dose reduction (ATV, COBI): underdosing?

✓ **Asunaprevir, 3D –drugs, MK 5172 and 8742**

- Lack of DDI data with COBI, DTG, MVC and many not-HIV drugs
- PK variability (demographics, genetics, clinical stages)

ID Unit

Andrea Calcagno

Letizia Marinaro

Laura Trentini

Cristina Tettoni

Chiara Alcantarini

Stefani Raviolo

Micol Ferrara

Maurizio Milesi

Chiara Montrucchio

Nicole Pagani

Prof Giovanni Di Perri



PK lab

Antonio D'Avolio

Jessica Cusato

Marco Simiele

Lorena Baietto

Alessandra Arialdo

Amedeo De Nicolò

Sarah Allegra

Debora Pensi

Mauro Sciandra

Viro lab

Tiziano Allice

Maria Grazia Milia

Valeria Ghisetti