

**10<sup>TH</sup> RESIDENTIAL COURSE ON  
CLINICAL PHARMACOLOGY OF ANTIRETROVIRALS**

# **TDM In Clinical Practice**

**LETIZIA MARINARO**


**Dip Malattie Infettive  
Università di Torino  
Ospedale Amedeo Di Savoia**

# Therapeutic Drug Monitoring (Tdm) For Antiretroviral Agents


Rationale for TDM of antiretrovirals points on two **key features**:

- ✓ Relationship between drug concentration and immunovirological benefit and in some cases, toxicity

Desired  
concentrations  
range



- ✓ Available clinical data show significant interpatient variability at equal dose intake



# DHHS Guidelines 2014

## TDM PROs

- Data showing that considerable **interpatient variability** in drug concentrations exists among patients who take the same dose
- Data indicating that relationships exist between the **concentration of drug** in the body and **anti-HIV effect** and, in some cases, **toxicities**
- Data from small prospective studies demonstrating that TDM **improved virologic response** and/or decreased the incidence of concentration-related drug toxicities .

# DHHS Guidelines 2014

## TDM CONs

- **lack of large prospective** studies demonstrating that TDM improves clinical and virologic outcomes
- **lack of established therapeutic** range of concentrations for all ARV drugs
- **inpatient variability** in ARV drug concentrations
- lack of widespread availability of clinical laboratories
- **shortage of experts** to assist with interpretation of ARV concentration data.

**Table 9a. Trough Concentrations of Antiretroviral Drugs for Patients Who Have Drug-Susceptible Virus**

Drug	Concentration (ng/mL)
<b>Suggested minimum target trough concentrations in patients with HIV-1 susceptible to the ARV drugs<sup>2-9</sup></b>	
Fosamprenavir (FPV)	400 (measured as amprenavir concentration)
Atazanavir (ATV)	150
Indinavir (IDV)	100
Lopinavir (LPV)	1000
Nelfinavir <sup>a</sup> (NFV)	800
Saquinavir (SQV)	100–250
Efavirenz (EFV)	1000
Nevirapine (NVP)	3000

<sup>a</sup> Measurable active (M8) metabolite

**Table 10b. Trough Concentrations of Antiretroviral Drugs for Treatment-Experienced Patients with Virologic Failure**

Drug	Concentration (ng/mL)
<b>Suggested minimum target trough concentrations for ART-experienced patients who have resistant HIV-1 strains</b>	
Maraviroc (MVC)	>50
Tipranavir (TPV)	20,500
<b>Median (Range) Trough Concentrations from Clinical Trials<sup>12-14</sup></b>	
Darunavir (DRV) (600 mg twice daily)	3300 (1255–7368)
Etravirine (ETR)	275 (81–2980)
Raltegravir (RAL)	72 (29–118)

# Scenarios for use of TDM

(DHHS 2014, Italian Guidelines 2014, BHIVA 2014)

- Clinically significant drug-drug or drug-food interactions
- Changes in pathophysiological states that may impair GI, hepatic, or renal function, thereby altering PK
- Persons such as pregnant women who may be at risk for virological failure as a result of their PK characteristics
- Treatment experienced patients
- Use of alternative dosing regimens
- Concentration-dependent toxicities
- Lack of expected virological response





**"We had a little trouble mixing your prescription.  
Can you stop back in 20 minutes?"**

## Risk of potentially clinically significant interactions

Study	Year	Setting	N	CSDI	Screening Tool	VL Effect
<i>de Maat</i>	2004	Netherlands (hospital)	115	26%	Liverpool website	N/A
<i>Shah et al</i>	2007	USA (Medicaid)	571	30%	Liverpool website; Micromedex	No VL impact
<i>Miller et al</i>	2007	USA (hospital)	153	41%	DHHS; PI; Micromedex	N/A
<i>Evans-Jones et al</i>	2010	UK (hospital)	159	27%	Liverpool website	N/A
<i>Marzolini et al</i>	2010	Switzerland (SHCS)	1497	40%	Liverpool website	No VL impact
<i>Kigen et al</i>	2011	Kenya (hospital)	996	34%	Liverpool website	N/A
<i>Patel et al</i>	2011	USA	190	34%	Lex-interact	N/A
<i>Cordova et al</i>	2013	Argentina	217	32%	Liverpool website	No VL impact
<i>Seden K et al</i>	2013	Uganda	2000	19%	Liverpool website	N/A


Shah et al. *CRUJ* 2007, Abstr 373  
 Miller et al. *Pharmacotherapy* 2007  
 Cordova E et al IAS 2013; MOPE031  
 Seden K et al; IAS 2013; MOPE035

Kigen et al. *Plos One* 2011  
 Patel Ann *Pharmacother* 2011  
 Marzolini et al. *Antivir Ther* 2010




# Rilpivirine and PPIs in HCV-coinfected pts

	AUC (ng h/mL)	Cmax (ng/mL)	Cmin (ng/mL)
<b>Patient A</b>	339	21	5
<b>Patient A without pantoprazole</b>	2562	124	90
<b>Patient B</b>	1901	113	39
<b>Patient C</b>	1518	128	34



# Rilpivirine and PPIs in HCV-coinfected pts

	AUC (ng h/mL)	C <sub>max</sub> (ng/mL)	C <sub>min</sub> (ng/mL)
<b>Patient A</b>	339	21	5
<b>Patient A without pantoprazole</b>	2562	124	90
<b>Patient B</b>	1901	113	39
<b>Patient C</b>	1518	128	34



**In two patients, the degree of hepatic impairment (Child B and C) counterbalanced the decrease in drug absorption**

*A. Calcagno, in press*

# DDIs and Pharmacogenetics

- Entity and “direction” of DDI determined by enzymes activity
- Interaction influenced by single nucleotide polymorphism (SNPs) of genes encoding for metabolizing enzymes

Calcagno et Al. Voriconazole and atazanavir: a CYP2C19-dependent manageable drug-drug interaction. *Pharmacogenomics*. 2014 Jul;15(10):1281-6.


- VORICONAZOLE metabolized by CYP2C19 (lesser extent by CYP2C9 and CYP3A4)
- RITONAVIR **INDUCES** CYP2C19 (inhibits 2C9 and 3A4)
- **Coadministration contraindicated** due 40% VOR Ctrough reduction

# DDIs and Pharmacogenetics

## Ritonavir and Voriconazole


*"Normal metabolizers"*

ATAZANAVIR  
RITONAVIR



# DDIs and Pharmacogenetics

## Ritonavir and Voriconazole




	1	1,5	2	2,5	3	3,5	4	5
<b>ATV</b>	553	196	251	352	228	110	139	155
<b>RTV</b>	273			447		229	65	90
<b>VOR</b>	5207	9921	9252	9673	8704	5984	1954	2357
<b>dose VOR</b>	<b>200x2</b>	<b>200x2</b>	<b>200x2</b>	<b>150x2</b>	<b>100x2</b>	<b>100x2</b>	<b>50x2</b>	<b>50x2</b>

# DDIs and Pharmacogenetics

## Ritonavir and voriconazole

*"Normal metabolizers"*

ATAZANAVIR  
RITONAVIR




↓ **VORICONAZOLE**  
(CYP2C19  
INDUCTION)

*"Poor and intermediate metabolizers"*

VORICONAZOLE

ATAZANAVIR  
RITONAVIR



↑ **VORICONAZOLE**  
(CYP3A4 INHIBITION)

# Italian HIV Guidelines 2014


- ✓ **Magnitude of a known interaction is unpredictable** in single patient, therefore TDM can be useful to check individual extent of such interaction [CIII].
- ✓ Net effect of multiple concomitant interactions is unpredictable, TDM is recommended [CIII].
- ✓ CLINICAL SIGNIFICANCE of an interaction could vary according to individual clinical variables, such as:
  - Genetic barrier of drugs involved
  - Immuno-virological status of the patient
  - Length of coadministration
- ✓ Combined evaluation of the latter with **TDM results** could allow the clinical management also of interactions otherwise considered at risk of therapeutical failure and/or toxicity



# MVC+COBI ?

- No data available

- Maraviroc 300 mg QD + Stribild  COBI boosting effect




	AUCss	Cmin	Cmax
	ng/mL*h	ng/mL	ng/mL
MVC 300 QD	4356	17	675
MEC (naive MERIT)		>25	

Unconventional regimen  
TDM at different timepoints  
confirms adequate plasma  
exposure



# Italian HIV Guidelines 2013

- ✓ **Magnitude of a known interaction is unpredictable** in single patient, therefore TDM can be useful to check individual extent of such interaction [CIII].
- ✓ **Net effect of multiple concomitant interactions is unpredictable, TDM is recommended** [CIII].
- ✓ CLINICAL SIGNIFICANCE of an interaction could vary according to individual clinical variables, such as:
  - Genetic barrier of drugs involved
  - Immuno-virological status of the patient
  - Length of coadministration
- ✓ Combined evaluation of the latter with **TDM results** could allow the clinical management also of interactions otherwise considered at risk of therapeutical failure and/or toxicity


# PROTEASE INHIBITORS + COBI ? STRIBILD + DRV 800



Cmin



AUC




Ramanathan S, et Al 13th International Workshop on Clinical Pharmacology of HIV Therapy [poster P08].

# PROTEASE INHIBITORS + COBI ? STRIBILD + DRV 800 QD

Patient with PHI

DRV,EVG,COBI plasma time-concentration profile



**Unconventional  
regimen**

**TDM confirms  
inadequate  
plasma exposure**

	T0	T1,5	T3	T4,5	T6	T8	T12	T24
DRV	334	3449	5405	6302	3996	2830	2491	287
EVG	68	426	812	1264	833	670	479	58
COBI	8	180	836	1092	640	380	141	10

# DOSE OPTIMIZATION



# CASE HISTORY

- Female, caucasian, 53 years old, HIV + since 1986, nadir CD4+ 4 cells/mL
- Starts ARV 1996:
  - 1996 AZT/3TC + IDV
  - 1998 D4T+3TC+ SQV
  - 1999 DDI+ D4T + EFV: DIZZINES
  - 2001 AZT/3TC + LPV/RTV: GI SYMPTOMS
  - 2004 AZT/3TC + EAP ATAZANAVIR/RTV improved GI SYMPTOMS
  - **2006 TDF/FTC + ATV/RTV ...HYPERBILIRUBINEMIA (3-4.5 mg/dL)**

**UNSTABLE THERAPY ACCEPTABILITY DUE TO MAJOR DEPRESSIVE DISORDER**

# ATV dose optimization

- **Low-Dose, Once-Daily Atazanavir/Ritonavir (200/100): An Effective Treatment for HIV Infected Patients in Thailand**  
Chetchotisakd P, JAIDS 2008
- **Simultaneous Population Pharmacokinetic Modelling of Atazanavir and Ritonavir in HIV-Infected Adults and Assessment of Different Dose Reduction Strategies**  
Schipani A, JAIDS 2013
- **A Maintenance Dose of Atazanavir/Ritonavir 200/100 mg Once Daily Is Effective in Virologically Suppressed HIV-1–Infected Patients**  
Lanzafame M, JAIDS 2013



# ATV dose optimization


## JAN 2012 : TDF/FTC + ATV/RTV

- HIV RNA not detectable, CD4+ 750 cells/ $\mu$ L, 35%
- BMI 20, hyperbilirubinemia 3.50 mg/mL, complains of scleral jaundice
- TDM : C trough ATV 799 ng/mL, RTV 59 ng/mL

## March 2012: ATV 200/RTV100+TDF/FTC

	PCR HIV	CD4+	BIL TOT	Ctrough ATV	Ctrough RTV
Jul 2012	<20	689	1.56		
Dec 2012	<20	730	0.63		
Jun 2013	Not detectable	917	1.88		
Dec 2013	Not detectable	650	2.01	345 ng/mL	76 ng/mL
May 2014	Not detectable	584	1.75		
Oct 2014	Not detectable	874	1.28	380 ng/mL	

## LASA trial: ongoing



Patients enrolled in Thailand. Maintenance trial, with primary analysis at Week 48 (HIV RNA suppression endpoint). Estimated Primary Completion Date: December 2014

# ATV dose optimization

- Reduced ATV dosing to 200 + RTV100: effective option in induction maintenance strategy
- Better compliance and tolerance due to reduced bilirubin levels
- Diminished risk of renal stones given the suggested correlation between stones development and ATV exposure
- Lowering dose could provide significant cost containment in resource-limited settings



“Frankly, darling, I think your doctor is a little obsessive about this compliance thing.”

# Drug schedule and polymedication

- 74-years old, diabetes, hypertension, IMA
- TDF/FTC + ATV/RTV, fully suppressed since 3 years, last VLs 230-350 copies/ml
- Insulin, aspirin, bisoprolol, rosuvastatin 10 mg, furosemide



# Drug schedule and polymedication

Doc: “What’s going on, dear patient, are you sure you’re always taking your pills as I told you?”

PT: “It’s true, doctor, I’m taking all my pills well!”

- TDM: ATV<sub>C12h</sub> 170 ng/ml (lower than expected)

Doc: “What’s going on, dear patient, are you sure you’re always taking your pills as I told you?”

PT: “I’m fully adherent, every single day I take Truvada and Norvir in the morning and Reyataz in the evening”

Doc: “.....!”


# Treatment failure & adherence

## TDM can:

- confirm poor adherence (low or undetectable)

## TDM cannot:

- confirm good adherence



Adequate concentration may only indicate patient has taken medications shortly before visit



# Challenges for TDM

- Knowledge of therapeutic ranges for ARVs (for wild type HIV and viruses with ↓ susceptibility)
- Widespread availability of quality controlled laboratories and timing of TDM results
- Interpretation and lack of dose adjustment strategies
- Lack of powered, prospective, controlled trials to demonstrate TDM improves outcome

# Future of TDM

- Evaluating TDM by RCT difficult: TDM must be integrated with all clinical factors
- ARVs are for life, on individual basis TDM can be a useful clinical tool
- Current drug levels for long term clinical response are not defined – studies warranted
- TDM-guided cost saving strategies

# Acknowledgments

**All the Staff of the Infectious Diseases Unit,  
University of Turin, Ospedale Amedeo Di Savoia**

**In particular:**

Prof. Giovanni Di Perri  
Prof. Stefano Bonora

**A special thanks to**

Doctor Andrea Calcagno  
for clinical data and  
slides kindly made  
available.

Chiara Alcantarini  
Sabrina Audagnotta  
Roberto Bertucci  
Lucio Boglione  
Paolo Bigliano  
Marino Bonasso  
Margherita Bracchi  
Chiara Cardellino  
Giuseppe Cariti  
Sinibaldo Carosella  
Silvia Corcione  
Cecilia Costa  
Micol Ferrara

Filippo Lipani  
Anna Lucchini  
Maurizio Milesi  
Chiara Montrucchio  
Simone Mornese  
Ilaria Motta  
Giancarlo Orofino  
Nicole Pagani  
Francesca Patti  
Stefania Raviolo  
Bernardino Salassa  
Silvia Scabini  
Maria Elisabetta Scarvaglieri  
Maria Laura Stella  
Maria Cristina Tettoni  
Alice Trentalange  
Laura Trentini

**All the Staff of the Clinical  
Pharmacological Laboratory**

**In particular:**

Antonio D'Avolio  
Sara Allegra  
Alessandra Ariaudo  
Lorena Baietto  
Jessica Gusato  
Amedeo De Nicolò  
Debora Pensi  
Marco Simiele  
Mauro Sciandra

**All the Staff of the Molecular  
Biology Laboratory**

**In particular:**

Valeria Ghisetti  
Tiziano Alice  
Tina Ruggiero  
Maria Grazia Milia